

Available online at www.sciencedirect.com

J. Math. Anal. Appl. 330 (2007) 245–258

Journal of
MATHEMATICAL
ANALYSIS AND
APPLICATIONS

www.elsevier.com/locate/jmaa

Dirichlet problem with discontinuous nonlinearities super-linear or asymptotically linear at infinity[☆]

Xian Wu^{*}, Tianjiu Leng

Department of Mathematics, Yunnan Normal University, Kunming, Yunnan 650092, PR China

Received 16 May 2006

Available online 24 August 2006

Submitted by S. Heikkilä

Abstract

In the present paper, a class of Dirichlet problem with discontinuous nonlinearities super-linear or asymptotically linear at infinity is studied, and some new existence theorems of solutions are obtained.

© 2006 Elsevier Inc. All rights reserved.

Keywords: Critical point; Locally Lipschitz function; Generalized gradient; (C) condition

1. Introduction

Throughout the paper, we assume $f : \bar{\Omega} \times R \rightarrow R$ be a locally bounded measurable function. Consider the following Dirichlet problem:

$$\begin{cases} -\Delta u = f(x, u), & \text{in } \Omega, \\ u = 0, & \text{on } \partial\Omega, \end{cases} \quad (1.1)$$

where $\Omega \subset R^N$ is a bounded smooth domain.

As $f(x, t) \in C(\bar{\Omega} \times R)$ is asymptotically linear, not super-linear, with respect to t at infinity, Zhou [8] researched the existence of positive solutions of problem (1.1) and obtained the following theorem.

[☆] This work is supported in partly the National Natural Science Foundation of China (10561011) and the Natural Science Foundation of Yunnan Province, China.

^{*} Corresponding author.

E-mail address: wuxian2001@yahoo.com.cn (X. Wu).

Theorem 1.1. *If*

- (H₁) $f(x, t) \in C(\bar{\Omega} \times R)$, $f(x, 0) = 0$, $\forall x \in \bar{\Omega}$, $f(x, t) \geq 0$, $\forall t \geq 0$, $x \in \bar{\Omega}$ and $f(x, t) \equiv 0$, $\forall t \leq 0$, $x \in \bar{\Omega}$;
 (H₂) for a.e. $x \in \Omega$, $\frac{f(x, t)}{t}$ is nondecreasing with respect to $t \geq 0$;
 (H₃) $\lim_{t \rightarrow 0} \frac{f(x, t)}{t} = p(x)$, $\lim_{t \rightarrow +\infty} \frac{f(x, t)}{t} = q(x) \neq 0$ uniformly in a.e. $x \in \Omega$, where $0 \leq p(x)$, $q(x) \in L^\infty(\Omega)$ and $\|p\|_\infty < \lambda_1$ ($\lambda_1 > 0$ is the first eigenvalue of $(-\Delta, H_0^1(\Omega))$), then

- (1) there is no positive solution to the problem (1.1) if $\Lambda > 1$;
- (2) the problem (1.1) has a positive solution if $\Lambda < 1$;
- (3) if $\Lambda = 1$, then problem (1.1) has a positive solution $u \in H_0^1(\Omega)$ if and only if there exists a constant $c > 0$ such that $u(x) = c\varphi_\Lambda(x)$ and $f(x, u) = q(x)u$ a.e. in $x \in \Omega$, where $\varphi_\Lambda(x) > 0$ is the function which achieves $\Lambda = \inf\{\int_\Omega |\nabla u|^2 dx : u \in H_0^1(\Omega), \int_\Omega q(x)u^2 dx = 1\}$ (see also [4]).

As $f(x, t) \in C(\bar{\Omega} \times R)$ is super-linear with respect to t at infinity, Zhou [8] obtained the following theorem.

Theorem 1.2. *If (H₁), (H₂) and (H₃) with $q(x) = +\infty$ and $\lim_{t \rightarrow +\infty} \frac{f(x, t)}{t^{r-1}} = 0$ uniformly in $x \in \Omega$ for some $r \in (2, \frac{2N}{N-2})$ if $N > 2$, or $r \in (2, +\infty)$ if $N = 1, 2$, then problem (1.1) has at least one positive solution $u \in H_0^1(\Omega)$.*

In 1976, Clarke [2] introduced the generalized gradients. In 1981, Chang [3] studied variational methods for nondifferentiable functionals by the generalized gradients and obtained several critical point theorems. As applications of these critical point theorems, Chang [3] researched partial differential equations with discontinuous nonlinearities. Since then, the results of Chang [3] become basic tools for the study of differential equation with discontinuous nonlinearities. Recently, Kyritsi and Papageorgiou [5] and Wu [7] studies critical points theory of nonsmooth function under nonsmooth C-condition, Bonanno and Giovannelli [1] researched nonsmooth eigenvalue problems and Marano and Motreanu [6] researched nonsmooth nonlinear boundary value problems under nonsmooth P.S.-condition. And as applications, Kyritsi and Papageorgiou [5] and Wu [7] researched the existence of solutions for hemivariational inequalities.

In the present paper, our main purpose is to study the existence of positive solution for problem (1.1) with $f: \bar{\Omega} \times R \rightarrow R$ be a locally bounded measurable function. Our main results contain Theorems 1.1 and 1.2 as their special cases and are different from these results in [1,5–7].

2. Preliminaries

Let X be a real Banach space and X^* the dual space of X . We denote by $\langle \cdot, \cdot \rangle$ the duality pairing between X^* and X . A functional $J: X \rightarrow R$ is called locally Lipschitz if for each $u \in X$ there exist a neighborhood U of u and a constant $L \geq 0$ such that

$$|J(v) - J(w)| \leq L\|v - w\|, \quad \forall v, w \in U.$$

For any $u, v \in X$, we define the generalized directional derivative $J^0(u; v)$ of J at point u along the direction v as

$$J^0(u; v) = \overline{\lim_{h \rightarrow 0, \lambda \downarrow 0}} \frac{1}{\lambda} [J(u + h + \lambda v) - J(u + h)].$$

The generalized gradient of the function J at u , denoted by $\partial J(u)$, is the set

$$\partial J(u) = \{w \in X^*: \langle w, v \rangle \leq J^0(u; v), \forall v \in X\}.$$

Set

$$\lambda(u) = \min_{w \in \partial J(u)} \|w\|.$$

A point $x_0 \in X$ is said to be a critical point of J if zero element in X^* $\theta \in \partial J(x_0)$. See [2,3] for the properties of the generalized directional derivative and the generalized gradient.

We need the following concept, which can be find in [5].

Definition 2.1. Let X be a normed linear space and $f: X \rightarrow R$ a locally Lipschitz function. We say that f satisfies the $(C)_c$ condition, if any sequence $\{x_n\} \subset X$ along which $f(x_n) \rightarrow c$ and $(1 + \|x_n\|)\lambda(x_n) \rightarrow 0$ possesses a convergent subsequence.

The following theorem is our main tool, which is Theorem 2.5 in [7].

Theorem 2.1. Let X be a reflexive Banach space and $f: X \rightarrow R$ a locally Lipschitz function. Assume that there exist a neighborhood U of 0, a point $x_0 \notin U$ and a constant β such that

$$f(0), \quad f(x_0) < \beta, \quad f|_{\partial U} \geq \beta.$$

Let $\Gamma = \{\varphi \in C([0, 1], X): \varphi(0) = 0, \varphi(1) = x_0\}$ and $c = \inf_{\varphi \in \Gamma} \max_{t \in [0, 1]} f(\varphi(t))$. Then $c \geq \beta$ and there exists a sequence $\{x_n\} \subset X$ such that $f(x_n) \rightarrow c$ and

$$(1 + \|x_n\|)\lambda(x_n) \rightarrow 0.$$

Furthermore, if f satisfies the $(C)_c$ condition, then c is a critical value of f .

3. Main results

Let $\Omega \subset R^N$ be a nonempty bounded open subset with a smooth boundary $\partial\Omega$. Given one locally bounded measurable function $f: \bar{\Omega} \times R \rightarrow R$. f is called superposition measurable if $u: \Omega \rightarrow R$ is measurable implies that $x \mapsto f(x, u(x))$ is measurable. Set

$$f^-(x, t) = \lim_{\delta \rightarrow 0} \inf_{|\xi - t| < \delta} f(x, \xi), \quad f^+(x, t) = \lim_{\delta \rightarrow 0} \sup_{|\xi - t| < \delta} f(x, \xi).$$

Then $f^-(x, \cdot)$ and $f^+(x, \cdot)$ are, respectively, lower semi-continuous and upper semi-continuous. Consider the following problems:

Find $u \in H_0^1(\Omega)$ such that

$$\begin{cases} -\Delta u \in [f^-(x, u(x)), f^+(x, u(x))], & \text{in } \Omega, \\ u = 0, & \text{on } \partial\Omega \end{cases} \quad (P_1)$$

or

$$\begin{cases} -\Delta u = f(x, u(x)), & \text{in } \Omega, \\ u = 0, & \text{on } \partial\Omega, \end{cases} \quad (P_2)$$

where f satisfies the following conditions:

(A₁) $f(x, 0) \equiv 0$ for all $x \in \bar{\Omega}$ and

$$f(x, t) \begin{cases} \geq (\neq) 0, & \text{if } t \geq 0, \\ \equiv 0, & \text{if } t \leq 0; \end{cases}$$

(A₂) $\frac{f(x, t)}{t}$ is nondecreasing in $t \geq 0$ for a.e. $x \in \Omega$;

(A₃) $\lim_{t \rightarrow 0} \frac{f(x, t)}{t} = m(x)$, $\lim_{t \rightarrow \infty} \frac{f(x, t)}{t} = q(x) \neq 0$, where $\|m\|_\infty < \lambda_1$, λ_1 is the first eigenvalue of the eigenvalue problem

$$\begin{cases} -\Delta u = \lambda u, & \text{in } \Omega, \\ u = 0, & \text{on } \partial\Omega; \end{cases}$$

(A₄) the functions f^- and f^+ are superposition measurable;

(A₅) there exists a constant r with

$$r \in \begin{cases} (2, 2^*), & \text{if } N > 2, \\ (2, +\infty), & \text{if } N = 1, 2 \end{cases}$$

such that

$$\lim_{t \rightarrow +\infty} \frac{f(x, t)}{t^{r-1}} = 0$$

uniformly for all $x \in \Omega$;

(A₆) there exists a set $\Omega_f \subset \Omega$ with $|\Omega_f|$ (the measure of Ω_f) = 0 such that the set

$$D_f := \bigcup_{x \in (\Omega \setminus \Omega_f)} \{t \in \mathbb{R}: f(x, \cdot) \text{ is discontinuous at } t\}$$

has measure zero, and for almost every $x \in \Omega$ each $t \in D_f$ the condition $f^-(x, t) \leq 0 \leq f^+(x, t)$ implies $f(x, t) = 0$.

Let Λ is the first eigenvalue of the eigenvalue problem

$$\begin{cases} -\Delta u = \lambda q(x)u, & \text{in } \Omega, \\ u = 0, & \text{on } \partial\Omega. \end{cases}$$

Then $\Lambda > 0$ is simple, isolated, $\Lambda = \inf\{\int_\Omega |\nabla u|^2 dx : u \in H_0^1(\Omega) \text{ and } \int_\Omega q(x)|u|^2 dx = 1\}$ and the infimum Λ can be achieved at some $\varphi_\Lambda > 0$ with $\int_\Omega q(x)|\varphi_\Lambda|^2 dx = 1$ (see [4]).

Theorem 3.1. (i) If conditions (A₁) to (A₄) hold, $q(x) \in L^\infty(\Omega)$ and $\Lambda > 1$, then the problem (P₁) has no any positive solution in $H_0^1(\Omega)$.

(ii) If conditions (A₁), (A₃), (A₄) hold, $q(x) \in L^\infty(\Omega)$ and $\Lambda < 1$, then the problem (P₁) possesses at least one positive solution in $H_0^1(\Omega)$. If adds the condition (A₆), then the problem (P₂) possesses at least one positive solution in $H_0^1(\Omega)$.

(iii) If conditions (A₁) to (A₄) hold, $q(x) \in L^\infty(\Omega)$ and $\Lambda = 1$, then the problem (P₁) possesses one positive solution $u \in H_0^1(\Omega) \Leftrightarrow$ there exists a positive constant α such that $u = \alpha\varphi_\Lambda$ and $f^+(x, u(x)) = q(x)u(x)$ for a.e. $x \in \Omega$.

Proof. Let $X = H_0^1(\Omega)$, $Y = L^2(\Omega)$, and

$$J(u) = \frac{1}{2} \int_\Omega |\nabla u|^2 dx - \int_\Omega F(x, u(x)) dx, \quad \forall u \in L^2(\Omega),$$

where $F(x, t) = \int_0^t f(x, s) ds$. Then the Sobolev embedding $H_0^1(\Omega) \hookrightarrow L^2(\Omega)$ is compact and J is well defined and locally Lipschitz in X .

Step 1. There exist constants $\rho, \beta > 0$ such that $J(u) \geq \beta$ whenever $u \in H_0^1(\Omega)$ and $\|u\| = \rho$.

Indeed, taking $2 < r < 2^*$. For any $0 < \varepsilon < \lambda_1 - \|m\|_\infty$, by (A₁) and (A₃), there exists a negative function $\alpha(x) \in L^\infty(\Omega)$ such that

$$|f(x, t)| < (m(x) + \varepsilon)|t| + \alpha(x)|t|^{r-1}$$

for all $x \in \Omega$ and all $t \in \mathbb{R}$. Consequently,

$$\begin{aligned} J(u) &= \frac{1}{2}\|u\|^2 - \int_{\Omega} F(x, u) dx \\ &\geq \frac{1}{2}\left(1 - \frac{\|m\|_\infty + \varepsilon}{\lambda_1}\right)\|u\|^2 - \frac{1}{r} \int_{\Omega} \alpha(x)|u|^r dx \\ &\geq \frac{1}{2}\left(1 - \frac{\|m\|_\infty + \varepsilon}{\lambda_1}\right)\|u\|^2 - \frac{1}{r}\|\alpha\|_s \|u\|_{\frac{rs}{s-1}}^r \\ &\geq \frac{1}{2}\left(1 - \frac{\|m\|_\infty + \varepsilon}{\lambda_1}\right)\|u\|^2 - c_1 \|u\|^r. \end{aligned}$$

Since $r > 2$ and $\frac{1}{2}(1 - \frac{\|m\|_\infty + \varepsilon}{\lambda_1}) > 0$, there exists small $\rho > 0$ such that

$$J(u) \geq \frac{1}{4}\left(1 - \frac{\|m\|_\infty + \varepsilon}{\lambda_1}\right)\rho^2 := \beta$$

whenever $u \in H_0^1(\Omega)$ and $\|u\| = \rho$.

Step 2. If $\Lambda < 1$, then $\lim_{t \rightarrow +\infty} J(t\varphi_\Lambda) = -\infty$. If $q(x) = +\infty$, then $\lim_{t \rightarrow +\infty} J(t\varphi_1) = -\infty$, where $\varphi_1 > 0$ is the eigenfunction corresponding to λ_1 .

The proofs are similar to the proofs of Lemma 2.2(b) in [8].

Step 3. Proof of (i).

If $u \in H_0^1(\Omega)$ is a positive solution of the problem (P₁), then

$$\begin{aligned} \int_{\Omega} |\nabla u|^2 dx &\leq \int_{\Omega} f^+(x, u(x))u(x) dx = \int_{\Omega} \lim_{\delta \rightarrow 0} \sup_{|\xi - u(x)| < \delta} f(x, \xi)u(x) dx \\ &\leq \int_{\Omega} \lim_{\delta \rightarrow 0} \sup_{|\xi - u(x)| < \delta} q(x)|\xi|u(x) dx = \int_{\Omega} q(x)|u|^2 dx. \end{aligned}$$

This shows $\Lambda \leq 1$. Hence, if $\Lambda > 1$, then the problem (P₁) has no any positive solution in $H_0^1(\Omega)$.

Step 4. Proof of (ii).

Since $\Lambda < 1$, by the conclusion of Step 2, there exists large $t_0 > 0$ such that $\|t_0\varphi_\Lambda\| > \rho$ and $J(t_0\varphi_\Lambda) < 0$. Set $M = [0, 1]$, $M^* = \{0, 1\}$, $\gamma^*(t) = tt_0\varphi_\Lambda$, $\forall t \in M$, and

$$\Gamma = \{\gamma \in C(M, X): \gamma(0) = 0, \gamma(1) = t_0\varphi_\Lambda\}, \quad c = \inf_{\gamma \in \Gamma} \max_{t \in [0, 1]} J(\gamma(t)).$$

Then $c \geq \beta > 0$. Now, we prove J satisfies the $(C)_c$ condition. Indeed, if any sequence $\{u_n\} \subset H_0^1(\Omega)$ such that $J(u_n) \rightarrow c$ and $(1 + \|u_n\|)\lambda(u_n) \rightarrow 0$, we shall prove that $\{u_n\}$ has a convergent subsequence.

Since $\lambda(u_n) = \min_{x^* \in \partial J(u_n)} \|x^*\|_{X^*}$, there exists $x_n^* \in \partial J(u_n)$ such that $\lambda(u_n) = \|x_n^*\|$. Hence

$$|\langle x_n^*, u_n \rangle| \leq \lambda(u_n)(1 + \|u_n\|) \rightarrow 0.$$

Since $x_n^* \in \partial J(u_n)$, there exists $z_n(x) \in [f^-(x, u_n(x)), f^+(x, u_n(x))]$ such that

$$\langle x_n^*, \varphi \rangle = \int_{\Omega} \nabla u_n \cdot \nabla \varphi \, dx - \int_{\Omega} z_n(x) \varphi(x) \, dx, \quad \forall \varphi \in X,$$

and hence

$$\|u_n\|^2 - \int_{\Omega} z_n(x) u_n(x) \, dx \rightarrow 0. \quad (3.1)$$

We claim that the sequence $\{u_n\}$ is bounded in X . Otherwise, we can assume $\|u_n\| \rightarrow \infty$. Set $w_n = \frac{u_n}{\|u_n\|}$. Then $\{w_n\}$ is bounded in X . Since X is a reflexive Banach space, we can assume that there exists $w \in X$ such that

$$w_n \rightharpoonup w \quad \text{in } X, \quad w_n \rightarrow w \quad \text{in } Y, \quad \text{and} \quad w_n(x) \rightarrow w(x) \quad \text{for a.e. } x \in \Omega.$$

If $w = 0$, then, by (A_1) , (A_3) and (3.1),

$$\begin{aligned} 1 &= \lim_{n \rightarrow \infty} \frac{1}{\|u_n\|^2} \int_{\Omega} z_n(x) u_n(x) \, dx = \lim_{n \rightarrow \infty} \int_{\Omega} \frac{z_n}{u_n} w_n^2 \, dx \\ &\leq \lim_{n \rightarrow \infty} \int_{\Omega} \frac{f^+(x, u_n(x))}{u_n} w_n^2 \, dx \leq c_1 \lim_{n \rightarrow \infty} \int_{\Omega} w_n^2 \, dx = 0. \end{aligned}$$

This is a contradiction. Hence $w \neq 0$. Consequently, for a.e. $x \in \Omega$, $u_n(x) \rightarrow +\infty$ as $n \rightarrow \infty$, and hence $\lim_{n \rightarrow \infty} \frac{f(x, u_n(x))}{u_n(x)} = q(x)$ for a.e. $x \in \Omega$. Consequently, by (A_3) , we can prove

$$\lim_{n \rightarrow \infty} \frac{f^+(x, u_n(x))}{u_n(x)} = q(x), \quad \lim_{n \rightarrow \infty} \frac{f^-(x, u_n(x))}{u_n(x)} = q(x),$$

and so, $\lim_{n \rightarrow \infty} \frac{z_n(x)}{u_n(x)} = q(x)$. Set

$$p_n(x) = \begin{cases} 0, & \text{if } u_n(x) \leq 0, \\ \frac{z_n(x)}{u_n(x)}, & \text{if } u_n(x) > 0. \end{cases}$$

By (A_1) and (A_3) there exists $M > 0$ such that $0 \leq p_n(x) \leq M$. Then for each $\varphi \in X$, one has

$$\begin{aligned} \int_{\Omega} p_n(x) w_n(x) \varphi(x) \, dx &= \int_{\Omega} p_n(x) [w_n(x) - w(x)] \varphi(x) \, dx + \int_{\Omega} p_n(x) w(x) \varphi(x) \, dx \\ &\rightarrow \int_{\Omega} q(x) w(x) \varphi(x) \, dx. \end{aligned}$$

Moreover,

$$\left| \int_{\Omega} \nabla w_n \cdot \nabla \varphi \, dx - \int_{\Omega} p_n(x) w_n(x) \varphi(x) \, dx \right| = \frac{1}{\|u_n\|} |\langle x_n^*, \varphi \rangle| \leq \frac{1}{\|u_n\|} \lambda(u_n) \|\varphi\| \rightarrow 0.$$

Hence

$$\int_{\Omega} \nabla w \cdot \nabla \varphi \, dx - \int_{\Omega} q(x) w(x) \varphi(x) \, dx = 0.$$

This contradicts that $\Lambda < 1$. Therefore, $\{u_n\}$ is bounded in X . Consequently, as the proof ²⁰ of Theorem 4.3 in [3] we may prove that $\{u_n\}$ possess a convergent subsequence, and hence J satisfies the $(C)_c$ condition. By Theorem 2.1 we know that c is a critical value of J . Hence there exist $u \in H_0^1(\Omega)$ and $z \in [f^-(x, u), f^+(x, u)]$ such that $J(u) = c \geq \beta > 0$ and

$$\int_{\Omega} \nabla u \cdot \nabla \varphi \, dx = \int_{\Omega} z(x) \varphi(x) \, dx \geq 0$$

for all nonnegative functions $\varphi \in C_0^1(\Omega)$. Consequently, by weak maximum principle we know that $u > 0$, i.e. $u \in H_0^1(\Omega)$ is a positive solution of the problem (P_1) . If adds the condition (A_6) , then as the proof of corresponding parts of Theorem 3.1 in [1] or Theorem 4.2 in [6] we can prove that the problem (P_2) possesses at least one positive solution in $H_0^1(\Omega)$.

Step 5. Proof of (iii).

Since $\Lambda = 1$,

$$\int_{\Omega} \nabla \varphi_{\Lambda} \cdot \nabla \varphi \, dx - \int_{\Omega} q(x) \varphi_{\Lambda}(x) \varphi(x) \, dx = 0, \quad \forall \varphi \in H_0^1(\Omega).$$

If $u \in H_0^1(\Omega)$ is a positive solution of the problem (P_1) , then there exists $z \in [f^-(x, u), f^+(x, u)]$ such that

$$\int_{\Omega} \nabla u \cdot \nabla \varphi \, dx = \int_{\Omega} z(x) \varphi(x) \, dx, \quad \forall \varphi \in H_0^1(\Omega). \quad (3.2)$$

Hence

$$\int_{\Omega} \nabla u \cdot \nabla \varphi_{\Lambda} \, dx = \int_{\Omega} z(x) \varphi_{\Lambda}(x) \, dx,$$

and hence

$$\begin{aligned} \int_{\Omega} q(x) \varphi_{\Lambda}(x) u(x) \, dx &= \int_{\Omega} \nabla \varphi_{\Lambda} \cdot \nabla u \, dx = \int_{\Omega} z(x) \varphi_{\Lambda}(x) \, dx \\ &\leq \int_{\Omega} f^+(x, u) \varphi_{\Lambda}(x) \, dx \leq \int_{\Omega} q(x) u(x) \varphi_{\Lambda}(x) \, dx. \end{aligned}$$

Therefore,

$$\int_{\Omega} [q(x) u(x) - z(x)] \varphi_{\Lambda}(x) \, dx = \int_{\Omega} [f^+(x, u) - z(x)] \varphi_{\Lambda}(x) \, dx = 0.$$

By (A_2) we know $f^+(x, u) = z(x) = q(x)u(x)$ for a.e. $x \in \Omega$. Consequently, by (3.2) one has

$$\int_{\Omega} \nabla u \cdot \nabla \varphi \, dx = \int_{\Omega} q(x)u(x)\varphi(x) \, dx, \quad \forall \varphi \in H_0^1(\Omega).$$

This shows that $u > 0$ is an eigenfunction associated to Λ . Since Λ is simple, there exists a constant $\alpha > 0$ such that $u = \alpha\varphi_{\Lambda}$.

Conversely, if there exists a positive constant α such that $u = \alpha\varphi_{\Lambda}$ and $f^+(x, u(x)) = q(x)u(x)$ for a.e. $x \in \Omega$, then

$$\begin{aligned} \int_{\Omega} \nabla u \cdot \nabla \varphi \, dx &= \alpha \int_{\Omega} \nabla \varphi_{\Lambda} \cdot \nabla \varphi \, dx = \alpha \int_{\Omega} q(x)\varphi_{\Lambda}(x)\varphi(x) \, dx \\ &= \int_{\Omega} q(x)u(x)\varphi(x) \, dx = \int_{\Omega} f^+(x, u(x))\varphi(x) \, dx \end{aligned}$$

for all $\varphi \in H_0^1(\Omega)$. Hence $u = \alpha\varphi_{\Lambda}$ is a positive solution of the problem (P_1) . \square

Corollary 3.2. *Let conditions (A_1) to (A_4) hold and $q(x) \equiv l > 0$.*

- (i) *If $\lambda_1 > l$, then the problem (P_1) has no any positive solution in $H_0^1(\Omega)$;*
- (ii) *If $\lambda_1 < l < +\infty$, then the problem (P_1) possesses at least one positive solution in $H_0^1(\Omega)$;*
- (iii) *If $\lambda_1 = l$, then the problem (P_1) possesses one positive solution $u \in H_0^1(\Omega) \Leftrightarrow$ there exists a positive constant α such that $u = \alpha\varphi_1$ and $f^+(x, u(x)) = \lambda_1 u(x)$ for a.e. $x \in \Omega$.*

Proof. Notice that $\Lambda = \frac{\lambda_1}{l}$. The conclusion follows from Theorem 3.1. \square

Theorem 3.3. *Let conditions (A_1) – (A_5) hold and $q(x) \equiv +\infty$. If adds the condition (A_6) with*

$$\Omega_f := \bigcup_{t>0} \{x \in \Omega: f^+(x, t) > f^-(x, t)\},$$

then the problem (P_2) possesses at least one positive solution in $H_0^1(\Omega)$.

Proof. For any $0 < \varepsilon < \lambda_1 - \|m\|_{\infty}$, by (A_1) , (A_3) and that

$$\lim_{t \rightarrow +\infty} \frac{f(x, t)}{t^{r-1}} = 0$$

uniformly for all $x \in \Omega$, there exist a positive constant α such that

$$|f(x, t)| < (m(x) + \varepsilon)|t| + \alpha|t|^{r-1} \quad (3.3)$$

for all $x \in \Omega$ and all $t \in \mathbb{R}$. Hence there exist constants $\rho, \beta > 0$ such that $J(u) \geq \beta$ whenever $u \in H_0^1(\Omega)$ and $\|u\| = \rho$ (see Step 1 of the proof of Theorem 3.1).

Since $q(x) \equiv +\infty$, by the conclusion of Step 2 of the proof of Theorem 3.1, there exists large $t_0 > 0$ such that $\|t_0\varphi_1\| > \rho$ and $J(t_0\varphi_1) < 0$. Set $M = [0, 1]$, $M^* = \{0, 1\}$, $\gamma^*(t) = tt_0\varphi_1$, $\forall t \in M$, and

$$\Gamma = \{\gamma \in c(M, X): \gamma(0) = 0, \gamma(1) = t_0\varphi_1\}, \quad c = \inf_{\gamma \in \Gamma} \max_{t \in [0, 1]} J(\gamma(t)).$$

Then $c \geq \beta > 0$. Now, we prove J satisfies the $(C)_c$ condition. Indeed, if any sequence $\{u_n\} \subset H_0^1(\Omega)$ such that $J(u_n) \rightarrow c$ and $(1 + \|u_n\|)\lambda(u_n) \rightarrow 0$, we shall prove that $\{u_n\}$ has a convergent subsequence.

We claim that the sequence $\{u_n\}$ is bounded in X . Otherwise, we can assume $\|u_n\| \rightarrow \infty$. Set $w_n = \frac{u_n}{\|u_n\|}$. Then $\{w_n\}$, $\{w_n^+\}$ and $\{w_n^-\}$ are bounded in X . Since X is a reflexive Banach space, we can assume that there exists $w \in X$ such that

$$w_n \rightharpoonup w \quad \text{in } X, \quad w_n \rightarrow w \quad \text{in } Y, \quad \text{and} \quad w_n(x) \rightarrow w(x) \quad \text{for a.e. } x \in \Omega,$$

and

$$w_n^+ \rightharpoonup w^+ \quad \text{in } X, \quad w_n^+ \rightarrow w^+ \quad \text{in } Y, \quad \text{and} \quad w_n^+(x) \rightarrow w^+(x) \quad \text{for a.e. } x \in \Omega.$$

Set $\Omega_1 = \{x \in \Omega : w^+(x) > 0\}$. Since $q(x) = +\infty$, by (A_1) , (A_3) and (3.1),

$$\begin{aligned} 1 &= \lim_{n \rightarrow \infty} \frac{1}{\|u_n\|^2} \int_{\Omega} z_n(x) u_n(x) dx = \lim_{n \rightarrow \infty} \int_{\Omega} \frac{z_n}{u_n} w_n^2 dx \\ &\geq \lim_{n \rightarrow \infty} \int_{\Omega} \frac{f^-(x, u_n^+(x))}{u_n^+} (w_n^+)^2 dx \geq \lim_{n \rightarrow \infty} \int_{\Omega_1} \frac{f^-(x, u_n^+(x))}{u_n^+} (w_n^+)^2 dx = +\infty, \end{aligned}$$

whenever the measure of Ω_1 , $|\Omega_1| > 0$. This is a contradiction. Hence $|\Omega_1| = 0$. Consequently, $w^+(x) = 0$ for a.e. $x \in \Omega$, and so that, by (3.3) and $r \in (2, 2^*)$,

$$\lim_{n \rightarrow +\infty} \int_{\Omega} F(x, 2\sqrt{c} w_n^+(x)) dx = 0.$$

Consequently,

$$\lim_{n \rightarrow +\infty} J(2\sqrt{c} w_n) = 2c.$$

Let $g(u) = \int_{\Omega} F(x, u(x)) dx$ and $x_n^* \in \partial J(u_n)$ such that $\lambda(u_n) = \|x_n^*\|$. Then there exists $z_n \in \partial g(u_n)$ such that

$$\langle x_n^*, \varphi \rangle = \int_{\Omega} \nabla u_n \cdot \nabla \varphi dx - \langle z_n, \varphi \rangle, \quad \forall \varphi \in X.$$

Since

$$|\langle x_n^*, u_n \rangle| \leq \lambda(u_n)(1 + \|u_n\|) \rightarrow 0, \quad \|u_n\|^2 - \langle z_n, u_n \rangle \rightarrow 0.$$

Hence we may assume that

$$-\frac{1}{n} < \|u_n\|^2 - \langle z_n, u_n \rangle < \frac{1}{n}, \quad \forall n \geq 1. \quad (3.4)$$

For any $t \geq 0$,

$$\begin{aligned} J(tu_n) &= \frac{1}{2} t^2 \|u_n\|^2 - \int_{\Omega} F(x, tu_n(x)) dx \\ &< \frac{1}{2} t^2 \left[\frac{1}{n} + \langle z_n, u_n \rangle \right] - \int_{\Omega} F(x, tu_n(x)) dx \end{aligned}$$

$$\begin{aligned}
&= \frac{t^2}{2n} + \frac{1}{2} t^2 \langle z_n, u_n \rangle - \int_{\Omega} F(x, tu_n(x)) dx \\
&\leq \frac{t^2}{2n} + \frac{1}{2} t^2 g^0(u_n; u_n) - \int_{\Omega} F(x, tu_n(x)) dx \\
&\leq \frac{t^2}{2n} + \int_{\Omega} \left[\frac{1}{2} t^2 F^0(x, u_n(x); u_n(x)) - F(x, tu_n(x)) \right] dx \\
&= \frac{t^2}{2n} + \int_{u_n(x) > 0} \left[\frac{1}{2} t^2 u_n(x) f^+(x, u_n(x)) - F(x, tu_n(x)) \right] dx.
\end{aligned}$$

For any fixed $x \in \Omega$ and positive integer n , set

$$h(t) = \frac{1}{2} t^2 u_n^+(x) f^+(x, u_n^+(x)) - F(x, tu_n^+(x)).$$

Then the function $h(t)$ is absolutely continuous in any closed interval $[a, b] \subset [0, +\infty)$ and differentiable almost everywhere in $(0, +\infty)$ and for a.e. $t \in [0, +\infty)$, one has

$$\begin{aligned}
\frac{d}{dt} h(t) &= t u_n^+(x) f^+(x, u_n^+(x)) - f(x, tu_n^+(x)) u_n^+(x) \\
&= t \left[u_n^+(x) f^+(x, u_n^+(x)) - \frac{f(x, tu_n^+(x))}{t} u_n^+(x) \right].
\end{aligned}$$

Consequently, as $0 \leq t_1 \leq 1$, one has

$$\begin{aligned}
h(1) - h(t_1) &= \int_{t_1}^1 \frac{d}{dt} h(t) dt \\
&= \frac{1}{2} (1 - t_1^2) u_n^+(x) f^+(x, u_n^+(x)) - \int_{t_1}^1 f(x, tu_n^+(x)) u_n^+(x) dt \\
&\geq \frac{1}{2} (1 - t_1^2) u_n^+(x) [f^+(x, u_n^+(x)) - f(x, u_n^+(x))] \\
&\geq 0,
\end{aligned}$$

i.e. $h(t_1) \leq h(1)$ for all $t_1 \in [0, 1]$. Consequently, for all $t \in [0, 1]$, one has

$$J(tu_n) \leq \frac{t^2}{2n} + \int_{u_n(x) > 0} \left[\frac{1}{2} u_n(x) f^+(x, u_n(x)) - F(x, u_n(x)) \right] dx.$$

On the other hand,

$$\begin{aligned}
J(u_n) &= \frac{1}{2} \|u_n\|^2 - \int_{\Omega} F(x, u_n(x)) dx \\
&> \frac{1}{2} \left[-\frac{1}{n} + \langle z_n, u_n \rangle \right] - \int_{\Omega} F(x, u_n(x)) dx
\end{aligned}$$

$$\begin{aligned}
&= -\frac{1}{2n} - \frac{1}{2} \langle z_n, -u_n \rangle - \int_{\Omega} F(x, u_n(x)) dx \\
&\geq -\frac{1}{2n} - \frac{1}{2} g^0(u_n; -u_n) - \int_{\Omega} F(x, u_n(x)) dx \\
&\geq -\frac{1}{2n} - \frac{1}{2} \int_{\Omega} F^0(x, u_n(x); -u_n(x)) dx - \int_{\Omega} F(x, u_n(x)) dx \\
&= -\frac{1}{2n} + \int_{u_n(x) > 0} \left[\frac{1}{2} u_n(x) f^-(x, u_n(x)) - F(x, u_n(x)) \right] dx.
\end{aligned}$$

Hence, for all $t \in [0, 1]$, one has

$$J(tu_n) \leq \frac{1+t^2}{2n} + \int_{u_n(x) > 0} \left[\frac{1}{2} u_n(x) [f^+(x, u_n(x)) - f^-(x, u_n(x))] \right] dx + J(u_n).$$

Since $|\Omega_f| = 0$,

$$\int_{u_n(x) > 0} \left[\frac{1}{2} u_n(x) [f^+(x, u_n(x)) - f^-(x, u_n(x))] \right] dx = 0.$$

Hence

$$J(tu_n) \leq \frac{1+t^2}{2n} + J(u_n).$$

Consequently,

$$J(2\sqrt{c}w_n) = J\left(\frac{2\sqrt{c}}{\|u_n\|} u_n\right) \leq \frac{1}{2n} \left(1 + \frac{4c}{\|u_n\|^2}\right) + J(u_n)$$

and hence $2c \leq c$. This contradicts that $c > 0$. Hence the sequence $\{u_n\}$ is bounded in X . Consequently, as the proof 2^0 of Theorem 4.3 in [3] we may prove that $\{u_n\}$ possess a convergent subsequence, and hence J satisfies the $(C)_c$ condition. By Theorem 2.1 we know that c is a critical value of J . Hence there exist $u \in H_0^1(\Omega)$ and $z \in [f^-(x, u), f^+(x, u)]$ such that $J(u) = c \geq \beta > 0$ and

$$\int_{\Omega} \nabla u \cdot \nabla \varphi dx = \int_{\Omega} z(x) \varphi(x) dx \geq 0$$

for all nonnegative functions $\varphi \in C_0^1(\Omega)$. Consequently, by weak maximum principle we know that $u > 0$, i.e. $u \in H_0^1(\Omega)$ is a positive solution of the problem (P_1) . The rest of the proof is the same with Theorem 3.1. \square

Remark. Theorems 3.1 and 3.3 generalize and improve Theorems 1.1 and 1.2 to the case that $f: \bar{\Omega} \times R \rightarrow R$ is only locally bounded measurable function, respectively. Moreover, by Theorems 3.1 and 3.3 we know that the condition $f(x, t) \in C(\bar{\Omega} \times R)$ in Theorems 1.1 and 1.2 may be weakened as $f(x, t)$ is a Caratheodory function.

The following Theorem 3.4 and Corollary 3.5 are new results, too.

Theorem 3.4. Let conditions (A_1) , (A_3) – (A_5) hold and $q(x) \equiv +\infty$. If there exists $\mu \geq r$ such that

$$\liminf_{t \rightarrow +\infty} \frac{tf(x, t) - 2F(x, t)}{|t|^\mu} > 0,$$

then the problem (P_1) possesses at least one positive solution in $H_0^1(\Omega)$. If adds the condition (A_6) , then the problem (P_2) possesses at least one positive solution in $H_0^1(\Omega)$.

Proof. It is sufficient to prove that J satisfies the $(C)_c$ condition from the proof of Theorem 3.3. By (A_1) , (A_3) and (A_5) , there exists a constant $c_1 > 0$ such that

$$0 \leq f(x, t) \leq c_1(1 + |t|^{r-1}), \quad \forall t \in \mathbb{R}.$$

Hence

$$F(x, t) = \int_0^t f(x, s) ds \leq c_1 + c_2 t^r, \quad \forall t \geq 0. \quad (3.5)$$

By

$$\liminf_{t \rightarrow +\infty} \frac{tf(x, t) - 2F(x, t)}{|t|^\mu} > 0$$

we can prove that

$$\liminf_{t \rightarrow +\infty} \frac{tf^-(x, t) - 2F(x, t)}{t^\mu} > 0. \quad (3.6)$$

Indeed, there exist $\varepsilon_0 > 0$, $M_0 > 0$ such that

$$\frac{sf(x, s) - 2F(x, s)}{s^\mu} > \varepsilon_0, \quad \forall s \geq M_0.$$

For each fixed t with $t > M_0 + 2$, as $\delta \in (0, 1)$ is enough small, one has $\frac{t}{\xi} > 0$ and

$$\xi > M_0 + 2, \quad \frac{t\xi^\mu}{t^\mu\xi} > \frac{1}{2}, \quad \frac{2F(x, \xi)}{t^{\mu-1}\xi} - \frac{2F(x, t)}{t^\mu} > -\frac{\varepsilon_0}{3}$$

for all ξ with $|\xi - t| < \delta$. Hence

$$\begin{aligned} \frac{tf(x, \xi) - 2F(x, t)}{t^\mu} &= \frac{\frac{t}{\xi} \cdot \xi f(x, \xi) - 2F(x, t)}{t^\mu} > \frac{\frac{t}{\xi} [2F(x, \xi) + \varepsilon_0 \xi^\mu] - 2F(x, t)}{t^\mu} \\ &= \frac{2F(x, \xi)}{t^{\mu-1}\xi} - \frac{2F(x, t)}{t^\mu} + \frac{t\xi^\mu}{t^\mu\xi} \varepsilon_0 > \frac{\varepsilon_0}{6}. \end{aligned}$$

Consequently,

$$\liminf_{t \rightarrow +\infty} \frac{tf^-(x, t) - 2F(x, t)}{t^\mu} > 0.$$

Assume $\{u_n\} \subset H_0^1(\Omega)$ such that $\{J(u_n)\}$ is bounded and $(1 + \|u_n\|)\lambda(u_n) \rightarrow 0$, where $\lambda(u_n) = \min_{w \in \partial J(u_n)} \|w\|$. Then there exists $M > 0$ such that

$$|J(u_n)| \leq M, \quad (1 + \|u_n\|)\lambda(u_n) \leq M.$$

Choose $w_n \in \partial J(u_n)$ such that $\|w_n\| = \lambda(u_n)$. Then there exists $v_n \in \partial g(u_n)$ such that

$$\langle w_n, v \rangle = \int_{\Omega} \nabla u_n \cdot \nabla v \, dx - \langle v_n, v \rangle, \quad \forall v \in H_0^1(\Omega).$$

Consequently, by (3.5)

$$\begin{aligned} \frac{1}{2} \|u_n\|^2 &= J(u_n) + \int_{\Omega} F(x, u_n(x)) \, dx \leq M + \int_{\Omega} [c_2(u_n^+(x))^r + c_1] \, dx \\ &\leq c_2 \int_{\Omega} (u_n^+(x))^r \, dx + c_3. \end{aligned} \quad (3.7)$$

Moreover, by (3.6) there exist constants $c_4 > 0$, $\delta_1 > 0$ such that

$$tf^-(x, t) - 2F(x, t) > c_4|t|^\mu, \quad \forall t \geq \delta_1.$$

By the locally boundedness of f there exist a constant $c_5 > 0$ such that

$$|tf^-(x, t) - 2F(x, t)| \leq c_5, \quad \forall 0 \leq t \leq \delta_1.$$

Hence

$$tf^-(x, t) - 2F(x, t) \geq c_4|t|^\mu - c_6, \quad \forall t \geq 0.$$

Let $g(u) = \int_{\Omega} F(x, u(x)) \, dx$. Consequently,

$$\begin{aligned} 3M &\geq 2J(u_n) - \langle w_n, u_n \rangle = -2 \int_{\Omega} F(x, u_n(x)) \, dx - \langle v_n, -u_n \rangle \\ &\geq -2 \int_{\Omega} F(x, u_n(x)) \, dx - g^0(u_n; -u_n) \\ &\geq -2 \int_{\Omega} F(x, u_n(x)) \, dx + \int_{u_n(x) < 0} u_n(x) f^+(x, u_n(x)) \, dx \\ &\quad + \int_{u_n(x) > 0} u_n(x) f^-(x, u_n(x)) \, dx \\ &= \int_{u_n(x) > 0} [u_n(x) f^-(x, u_n(x)) - 2F(x, u_n(x))] \, dx \\ &\geq \int_{\Omega} [c_4(u_n^+(x))^\mu - c_6] \, dx. \end{aligned}$$

Therefore, $\int_{\Omega} (u_n^+(x))^\mu \, dx$ is bounded. Since $\mu \geq r$, by (3.7), $\|u_n\|$ is bounded. The rest of the proof is the same with proof 2^o of Theorem 4.3 in [3]. \square

Corollary 3.5. Let $f: \bar{\Omega} \times \mathbb{R} \rightarrow \mathbb{R}$ be a locally bounded Caratheodory function and conditions (A₁), (A₃) and (A₅) hold with $q(x) \equiv +\infty$. If there exists $\mu \geq r$ such that

$$\liminf_{t \rightarrow +\infty} \frac{tf(x, t) - 2F(x, t)}{|t|^\mu} > 0,$$

then the problem (P₂) possesses at least one positive solution in $H_0^1(\Omega)$.

References

- [1] G. Bonanno, N. Giovannelli, An eigenvalue Dirichlet problem involving the p -Laplacian with discontinuous nonlinearities, *J. Math. Anal. Appl.* 308 (2005) 596–604.
- [2] F.H. Clarke, A new approach to Lagrange multipliers, *Math. Oper. Res.* 1 (1976) 165–174.
- [3] K.C. Chang, Variational methods for non-differentiable functionals and their applications to partial differential equations, *J. Math. Anal. Appl.* 80 (1981) 102–129.
- [4] M. Cuesta, Eigenvalue problems for the p -Laplacian with indefinite weights, *Electron. J. Differential Equations* 2001 (33) (2001) 1–9.
- [5] S.T. Kyrtsi, N.S. Papageorgiou, Nonsmooth critical point theory on closed convex sets and nonlinear hemivariational inequalities, *Nonlinear Anal.* 61 (2005) 373–403.
- [6] S. Marano, D. Motreanu, On a three critical points theorem for non-differentiable functions and applications to nonlinear boundary value problems, *Nonlinear Anal.* 48 (2002) 37–52.
- [7] Xina Wu, The study of critical point theory for non-differentiable functionals with applications to differential equations, *Nonlinear Anal.*, in press.
- [8] Huan-Song Zhou, An application of a Mountain Pass Theorem, *Acta Math. Sinica (English Ser.)* 18 (1) (2002) 27–36.